

SB23-144 Prescription Drugs for Chronic Pain

Sponsor: Senator Ginal

A recent study showed that as many as 40% of primary care physician offices will NOT see a new patient if they are using opiate pain medications.^{1 2 3}

Doctors' fear of increased scrutiny for prescribing medications for patients with a legitimate medical need has had a chilling effect on care.

Overly restrictive policies limiting controlled substances for pain to try to reduce overdose deaths has created another crisis:

Coloradans living with chronic pain are unable to access the treatment they and their doctors feel they need, resulting in increased disability, decreased function, suicides, and an increase in overdoses as pain patients try to treat their own pain.^{4 7 8}

Tapering a patient down or off opioid medication can actually increase their risk of overdose and suicide.^{6 10}

CDC Data suggest that illicit fentanyl and other illegally obtained controlled substances, tainted substances, or the use of multiple substances in combination drive overdose deaths.^{9 5}

This Bill:

→ Protects access to healthcare by prohibiting a healthcare provider or health system from refusing to treat a patient solely because of the patient's use of prescribed opioids or opioids at a particular dose.

→ Clarifies that a prescribing healthcare provider treating a patient for chronic pain is not subject to disciplinary action solely for prescribing a controlled substance at a dose above a set or predetermined dose threshold.

→ Prohibits a healthcare provider from forcing a patient to taper their medication below the dose they need, solely because of predetermined dose threshold recommendations, not for independent, clinical reasons.

→ Prohibits a pharmacy, health insurance carrier, or pharmacy benefit manager from refusing to fill or approve the coverage for a drug solely on the basis of the dosage requirement of a patient.

→ Improves access to controlled substances for patients with genuine medical need, including those in palliative, hospice, or end-of-life care. Policies around controlled substances for pain have left many patients struggling to access pain control and medications.

→ Respects that treatment decisions should be individualized and made between the doctor and the patient. Ensures that clinical discretion--whether by a pharmacist or physician--will take precedence over one-size-fits-all policies.

→ Helps protect Coloradans who need opioids for chronic pain management by improving access to care and enables them to get medications from a pharmacy of their choice.

This Bill Will NOT:

⊘ Force doctors to prescribe opiates, other controlled substances, or any other treatments.

⊘ Prevent people who are illegally diverting medications from being prosecuted.

⊘ Prohibit providers from referring patients who generally require care outside of their area of expertise.

[1] (Gavin, Michigan Medicine 2021)

[2] (Lagisetty, JAMA Netw Open 2019)

[3] (Quest Diagnostics survey, 2019)

[4] (Lagisetty, PAIN 2021)

[5] (Bitting, MMWR Morb Mortal Wkly Rep 2022)

[6] (Magnan, JAMA Netw Open 2023)

[7] (Larochelle, JAMA Netw Open 2022)

[8] (Glanz, JAMA Netw Open 2019)

[9] (Fentanyl Facts, 2022)

[10] (Neprash, J Gen Intern Med 2021)

[11] (Brown, J of Pain and Symptom Manage 2020)

Key Terms:

Tolerance

A condition that occurs when the body gets used to a medicine so that either more medicine or a different medicine may be needed.

Dependence

Physical adaptation to a medication so that the sudden absence of the agent causes withdrawal symptoms. Many medications other than controlled substances can cause dependence.

Addiction

A chronic, relapsing condition of compulsive drug use despite adverse consequences.

"MME"

Morphine Milligram Equivalent Dose - Used to convert the dose of an opiate to the equivalent dose of morphine in mg. Usually referred to as a daily total, IE 90 MME/day.



**MENTAL HEALTH
COLORADO**



**COLORADO
CROSS-DISABILITY
COALITION**

References:

- Agnoli, A., Xing, G., Tancredi, D. J., Magnan, E., Jerant, A., & Fenton, J. J. (2021). Association of Dose Tapering With Overdose or Mental Health Crisis Among Patients Prescribed Long-term Opioids. *JAMA*, 326(5), 411–419. <https://doi.org/10.1001/jama.2021.11013>
- Bitting, J. (2022). Notes from the Field: Overdose Deaths Involving Para-fluorofentanyl — United States, July 2020–June 2021. *MMWR. Morbidity and Mortality Weekly Report*, 71. <https://doi.org/10.15585/mmwr.mm7139a3>
- Brown, J. H., Torres, H. P., Maddi, R. D., Williams, J. L., Dibaj, S. S., Liu, D., & Bruera, E. (2020). Cancer Patients' Perceived Difficulties Filling Opioid Prescriptions After Receiving Outpatient Supportive Care. *Journal of Pain and Symptom Management*, 60(5), 915–922. <https://doi.org/10.1016/j.jpainsymman.2020.06.015>
- Fentanyl Facts. (2022, February 23). <https://www.cdc.gov/stopoverdose/fentanyl/index.html>
- Fenton, J. J., Magnan, E., Tseregounis, I. E., Xing, G., Agnoli, A. L., & Tancredi, D. J. (2022). Long-term Risk of Overdose or Mental Health Crisis After Opioid Dose Tapering. *JAMA Network Open*, 5(6), e2216726. <https://doi.org/10.1001/jamanetworkopen.2022.16726>
- Gavin, K. (2021, January 25). Pain Patients Who Take Opioids Can't Get in the Door at Half of Primary Care Clinics. *Michigan Medicine*. <https://www.michiganmedicine.org/health-lab/pain-patients-who-take-opioids-cant-get-door-half-primary-care-clinics>
- Glanz, J. M., Binswanger, I. A., Shetterly, S. M., Narwaney, K. J., & Xu, S. (2019). Association Between Opioid Dose Variability and Opioid Overdose Among Adults Prescribed Long-term Opioid Therapy. *JAMA Network Open*, 2(4), e192613. <https://doi.org/10.1001/jamanetworkopen.2019.2613>
- James, J. R., Scott, J. M., Klein, J. W., Jackson, S., McKinney, C., Novack, M., Chew, L., & Merrill, J. O. (2019). Mortality After Discontinuation of Primary Care-Based Chronic Opioid Therapy for Pain: A Retrospective Cohort Study. *Journal of General Internal Medicine*, 34(12), 2749–2755. <https://doi.org/10.1007/s11606-019-05301-2>
- Lagisetty, P. A., Healy, N., Garpestad, C., Jannausch, M., Tipirneni, R., & Bohnert, A. S. B. (2019). Access to Primary Care Clinics for Patients With Chronic Pain Receiving Opioids. *JAMA Network Open*, 2(7), e196928. <https://doi.org/10.1001/jamanetworkopen.2019.6928>
- Lagisetty, P., Macleod, C., Thomas, J., Slat, S., Kehne, A., Heisler, M., Bohnert, A. S. B., & Bohnert, K. M. (2021). Assessing reasons for decreased primary care access for individuals on prescribed opioids: An audit study. *PAIN*, 162(5), 1379. <https://doi.org/10.1097/j.pain.0000000000002145>
- Larochelle, M. R., Lodi, S., Yan, S., Clothier, B. A., Goldsmith, E. S., & Bohnert, A. S. B. (2022). Comparative Effectiveness of Opioid Tapering or Abrupt Discontinuation vs No Dosage Change for Opioid Overdose or Suicide for Patients Receiving Stable Long-term Opioid Therapy. *JAMA Network Open*, 5(8), e2226523. <https://doi.org/10.1001/jamanetworkopen.2022.26523>
- Mark, T. L., & Parish, W. (2019). Opioid medication discontinuation and risk of adverse opioid-related health care events. *Journal of Substance Abuse Treatment*, 103, 58–63. <https://doi.org/10.1016/j.jsat.2019.05.001>
- Nephrash, H. T., Gaye, M., & Barnett, M. L. (2021). Abrupt Discontinuation of Long-term Opioid Therapy Among Medicare Beneficiaries, 2012–2017. *Journal of General Internal Medicine*, 36(6). <https://doi.org/10.1007/s11606-020-06402-z>
- Oliva, E. M., Bowe, T., Manhapra, A., Kertesz, S., Hah, J. M., Henderson, P., Robinson, A., Paik, M., Sandbrink, F., Gordon, A. J., & Trafton, J. A. (2020). Associations between stopping prescriptions for opioids, length of opioid treatment, and overdose or suicide deaths in US veterans: Observational evaluation. *BMJ*, 368, m283. <https://doi.org/10.1136/bmj.m283>
- Perez, H. R., Buonora, M., Cunningham, C. O., Heo, M., & Starrels, J. L. (2020). Opioid Taper Is Associated with Subsequent Termination of Care: A Retrospective Cohort Study. *Journal of General Internal Medicine*, 35(1), 36–42. <https://doi.org/10.1007/s11606-019-05227-9>
- Quest Diagnostics—Health Trends Report. (2019). *Health Trends*.